

Conzol™

(Miconazole Nitrate)

Conzol is a synthetic antifungal agent that is indicated for the treatment of fungal infections in dogs and cats caused by *Microsporum canis*, *Microsporum gypseum* and *Trichophyton mentagrophytes*. Contains 1.15% miconazole nitrate (equivalent to 1% miconazole base by weight), polyethylene glycol 400 and ethyl alcohol 55%. Prescription.

ITEM#	DESCRIPTION	SIZE
501008	Conzol Lotion	60 mL
502008	Conzol Spray	120 mL
503008	Conzol Spray	240 mL



Available exclusively through your veterinarian from:



(800) 824-3703
www.mwivet.com



(888) 722-2242
www.aahamarketlink.com


Conzol™
(Miconazole Nitrate)**Topical Antifungal Agent****MICONAZOLE NITRATE**
CONZOL™ TOPICAL LOTION AND SPRAY

ANADA #200-196, Approved by FDA.
Veterinary
For Topical Use in Dogs and Cats Only.

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION: Conzol™ is a synthetic antifungal agent for use in dogs and cats. It contains: 1.15% miconazole nitrate (equivalent to 1% miconazole base by weight), polyethylene glycol 400 and ethyl alcohol 55%.

INDICATIONS: Conzol™ is indicated for the treatment of fungal infections in dogs and cats caused by *Microsporum canis*, *Microsporum gypseum* and *Trichophyton mentagrophytes*.

PRECAUTIONS: In the event of sensitization or irritation due to Conzol™, treatment should be discontinued. Avoid contact with eyes, since irritation may result. Wash hands thoroughly after administration to avoid spread of fungal infection.

DOSAGE AND ADMINISTRATION:

Accurate diagnosis of the infecting organism is essential. Identification should be made either by direct microscopic examination of a mounting of infected tissue in a solution of potassium hydroxide, or by culture on an appropriate medium.

Spray affected areas from a distance of 2 to 4 inches to apply a light covering, once daily for 2 to 4 weeks. Medication must be continued until the infecting organism is completely eradicated as indicated by appropriate clinical or laboratory examination. If no improvement is noticed within 2 weeks, diagnosis should be re-evaluated. Difficult cases may require treatment for 6 weeks.

General measures in regard to hygiene should be observed to control sources of infection or reinfection. Clipping of hair around and over the sites of infection should be done at the start of treatment and again as necessary.

HOW SUPPLIED: Conzol™ Lotion in 60mL container and Conzol™ Spray in 120 mL and 240 mL containers.