

DextroLyte™

Injection

This sterile solution of electrolytes and dextrose is for use in conditions associated with fluid and electrolyte loss, such as dehydration, shock, vomiting and diarrhea, particularly when an immediate source of energy is also indicated. For use in adult cattle and horses, calves, ponies, foals, adult sheep and swine. Administer by intravenous or intraperitoneal injection; do not administer intraperitoneally to horses. Prescription.

ITEM#	DESCRIPTION	SIZE
510183	DextroLyte Injection	500 mL
510184	DextroLyte Injection	1000 mL



Available exclusively through your veterinarian from:



(800) 824-3703
www.mwivet.com



(888) 722-2242
www.aahamarketlink.com

DEXTROLYTE™

Injection

Sterile Solution Electrolytes Dextrose

For animal use only.

Keep out of reach of children and pets.

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

CONTAINS: Each 500 mL of sterile aqueous solution contains:

Dextrose • H ₂ O	12.50 g
Sorbitol	12.50 g
Sodium Lactate	3.95 g
Sodium Chloride	2.40 g
Potassium Chloride	0.37 g
Magnesium Chloride • 6H ₂ O	0.21 g
Calcium Chloride • 2H ₂ O	0.19 g

Milliequivalents per liter

Cations

Sodium	153 mEq/L
Potassium	9 mEq/L
Calcium	6 mEq/L
Magnesium	4 mEq/L

Anions

Chloride	101 mEq/L
Lactate	71 mEq/L

Osmolarity (calc.): 602 mOsmol/L

CAUTION: This product contains no preservatives. Use entire contents when first opened. Discard any unused solution.

STORAGE: Store between 15°C-30°C (59°F-86°F).

INDICATIONS: For use in conditions associated with fluid and electrolyte loss, such as dehydration, shock, vomiting and diarrhea, particularly when an immediate source of energy is also indicated.

DOSAGE AND ADMINISTRATION: Warm solution to body temperature and administer slowly (10 to 30 mL per minute) by intravenous or intraperitoneal injection, using strict aseptic procedures.

Adult Cattle and Horses: 1000 to 2000 mL

Calves, Ponies and Foals: 500 to 1000 mL

Adult Sheep and Swine: 500 to 1000 mL

These are suggested dosages. The actual amount and rate of fluid administration must be judged by the veterinarian in relation to the condition being treated and the clinical response of the animal, being careful to avoid overhydration.

CONTRAINDICATIONS: Do not administer intraperitoneally to horses.

MADE IN USA

Distributed by: MWI, Boise, ID 83705

(888) 694-8381

www.vetone.net

Net Contents: NDC

500 mL - V1 510183

1000 mL - V1 510184

NAC No.: 13151460